arXiv:1304.5369v1 [math.CA] 19 Apr 2013

SOME NEW INEQUALITIES FOR TRIGONOMETRIC FUNCTIONS AND CORRESPONDING ONES FOR MEANS

ZHEN-HANG YANG

Abstract. In this paper, the versions of trigonometric functions of certain known inequalities for hyperbolic ones are proved, and then corresponding inequalities for means are presented.

1. Introduction

We begin with the following lemma.

Lemma 1. *Let $M(x,y)$ be a homogeneous mean of positive arguments x and y. Then*

$$M(x,y)=\sqrt{xy}M\left(e^{t},e^{-t}\right),$$

where $t=\frac{1}{2}\ln(x/y)$.

Using this lemma, almost all of inequalities for homogeneous symmetric bivariate means can be transformed equivalently into the corresponding ones for hyperbolic functions and vice versa. More specifically, let $L(x,y)$, $I(x,y)$ and $A_r(x,y)$ denote the logarithmic, identric and r-order power means of two positive real numbers x and y with $x\neq y$ defined by

$$\begin{aligned}L &= L(x,y)=\frac{x-y}{\ln x-\ln y},\quad I=I(x,y)=e^{-1}\left(\frac{x^{x}}{y^{y}}\right)^{1/(x-y)},\\ A_r &= A_r(x,y)=\left(\frac{x^{r}+y^{r}}{2}\right)^{1/r}\text{ if }r\neq 0\text{ and }G=A_0(x,y)=\sqrt{xy},\end{aligned}$$

respectively. Then we have

$$L\left(e^{t},e^{-t}\right)=\tfrac{\sinh t}{t},\ I\left(e^{t},e^{-t}\right)=e^{t\coth t-1},\ A_p\left(e^{t},e^{-t}\right)=\cosh^{1/p}pt,\ G\left(e^{t},e^{-t}\right)=1,$$

where $t=\frac{1}{2}\ln(x/y)>0$. And then, by Lemma 1, we can obtain some inequalities for hyperbolic functions from certain known ones for means mentioned previously. Here are some examples:

$$\begin{aligned}A_{2/3} &< I<A_{\ln 2}\Longrightarrow\\ \left(\cosh\frac{2t}{3}\right)^{3/2} &< e^{t\coth t-1}<\left(\cosh\left(t\ln 2\right)\right)^{1/\ln 2}\quad\text{(see [27], [20])}\end{aligned}\tag{1.1}$$

Date: April 15, 2013.
2010 *Mathematics Subject Classification.* Primary 26D05, 33B10; Secondary 26E60, 26D15 .
Key words and phrases. Trigonometric function, inequality, mean.
This paper is in final form and no version of it will be submitted for publication elsewhere.

$$\begin{aligned} A_{2/3} &< I<\sqrt{8}e^{-1}A_{2/3}\Longrightarrow \\ \left(\cosh\frac{2t}{3}\right)^{3/2} &< e^{t\coth t-1}<\sqrt{8}e^{-1}\left(\cosh\frac{2t}{3}\right)^{3/2}\quad \text{(see [30])} \end{aligned}\tag{1.2}$$

$$\sqrt{IG}<L<\frac{I+G}{2}\Longrightarrow\sqrt{e^{t\coth t-1}}<\frac{\sinh t}{t}<\frac{e^{t\coth t-1}+1}{2}\quad\text{(see [1])},\tag{1.3}$$

$$I>\frac{L+A}{2}\Longrightarrow e^{t\coth t-1}>\frac{\frac{\sinh t}{t}+\cosh t}{2}\quad\text{(see [21])}\tag{1.4}$$

$$\begin{aligned} \sqrt{AG} &< \sqrt{LI}<\frac{L+I}{2}<\frac{A+G}{2}\Longrightarrow \\ \sqrt{\cosh t} &< \sqrt{\tfrac{\sinh t}{t}e^{t\coth t-1}}<\tfrac{\frac{\sinh t}{t}+e^{t\coth t-1}}{2}<\tfrac{\cosh t+1}{2}\quad\text{(see [2])} \end{aligned}\tag{1.5}$$

$$\begin{aligned} \alpha A+(1-\alpha)G &< I<\beta A+(1-\beta)\,G\Longrightarrow \\ \alpha\cosh t+(1-\alpha) &< e^{t\coth t-1}<\beta\cosh t+(1-\beta)\quad\text{(see [3])} \end{aligned}\tag{1.6}$$

hold if and only if $\alpha\le 2/3$ and $\beta\ge 2/e$.

$$\begin{aligned} \left(\tfrac{2}{3}A^p+\tfrac{1}{3}G^p\right)^{1/p} &< I<\left(\tfrac{2}{3}A^q+\tfrac{1}{3}G^q\right)^{1/q}\Longrightarrow \\ \left(\tfrac{2}{3}\cosh^p t+\tfrac{1}{3}\right)^{1/p} &< e^{t\coth t-1}<\left(\tfrac{2}{3}\cosh^q t+\tfrac{1}{3}\right)^{1/q}\quad\text{(see [14])} \end{aligned}\tag{1.7}$$

hold if and only if $p\le 6/5$ and $q\ge(\log 3-\log 2)\,/\,(1-\log 2)$.

The aim of this paper is to give the versions of trigonometric functions listed above and others. The rest of this paper is organized as follows. In Section 2, we give some useful lemmas. Main results are contained in Section 3. In the last section, some inequalities for means corresponding to main results are given.

2. LEMMAS

Lemma 2 ([28], [4])**.** *Let $f,g:[a,b]\to\mathbb{R}$ be two continuous functions which are differentiable on (a,b). Further, let $g'\ne 0$ on (a,b). If f'/g' is increasing (or decreasing) on (a,b), then so are the functions*

$$t\mapsto\frac{f(t)-f(b)}{g(t)-g(b)}\quad\text{and}\quad t\mapsto\frac{f(t)-f(a)}{g(t)-g(a)}.$$

Lemma 3 ([5])**.** *Let a_n and b_n $(n=0,1,2,...)$ be real numbers and let the power series $A(t)=\sum_{n=1}^{\infty}a_nt^n$ and $B(t)=\sum_{n=1}^{\infty}b_nt^n$ be convergent for $|t|<R$. If $b_n>0$ for $n=0,1,2,...$, and a_n/b_n is strictly increasing (or decreasing) for $n=0,1,2,...$, then the function $A(t)/B(t)$ is strictly increasing (or decreasing) on $(0,R)$.*

Lemma 4 ([12, pp.227-229])**.** *We have*

$$\frac{1}{\sin t} = \frac{1}{t}+\sum_{n=1}^{\infty}\frac{2^{2n}-2}{(2n)!}|B_{2n}|t^{2n-1},\ |t|<\pi \tag{2.1}$$

$$\cot t = \frac{1}{t}-\sum_{n=1}^{\infty}\frac{2^{2n}}{(2n)!}|B_{2n}|t^{2n-1},\ \ |t|<\pi, \tag{2.2}$$

$$\tan t = \sum_{n=1}^{\infty}\frac{2^{2n}-1}{(2n)!}2^{2n}|B_{2n}|t^{2n-1},\ \ |t|<\pi/2, \tag{2.3}$$

$$\frac{1}{\sin^2 t} = \frac{1}{t^2}+\sum_{n=1}^{\infty}\frac{(2n-1)\,2^{2n}}{(2n)!}|B_{2n}|t^{2n-2},\ \ |t|<\pi, \tag{2.4}$$

where B_n is the Bernoulli number.

Lemma 5. *For every $t\in(0,\pi/2)$, $p\in(0,1]$, the function F_p defined by*

$$F_p\left(t\right)=\frac{t\cot t-1}{\ln\cos pt}$$

is increasing if $p\in(0,1/2]$ and decreasing if $p\in[\sqrt{10}/5,1]$. Consequently, for $p\in(0,1/2]$ we have

$$\frac{2}{3p^2}<\frac{t\cot t-1}{\ln\cos pt}<-\frac{1}{\ln\left(\cos\left(\pi p/2\right)\right)}. \tag{2.5}$$

It is reversed if $p\in[\sqrt{10}/5,1]$.

Proof. For $t\in(0,\pi/2)$, we define $f_1\left(t\right)=t\cot t-1$ and $f_2\left(t\right)=\ln\left(\cos pt\right)$, where $p\in(0,1]$. Note that $f_1\left(0^+\right)=f_2\left(0^+\right)=0$, then $F_p\left(t\right)$ can be written as

$$F_p\left(t\right)=\frac{f_1\left(t\right)-f_1\left(0^+\right)}{f_2\left(t\right)-f_2\left(0^+\right)}.$$

Differentiation and using (2.1) and (2.2) yield

$$\begin{aligned}\frac{f_1'\left(t\right)}{f_2'\left(t\right)} &= \frac{\frac{t}{\sin^2 t}-\cot t}{p\tan pt}=\frac{t\left(\frac{1}{t^2}+\sum_{n=1}^{\infty}\frac{(2n-1)2^{2n}}{(2n)!}|B_{2n}|t^{2n-2}\right)-\left(\frac{1}{t}-\sum_{n=1}^{\infty}\frac{2^{2n}}{(2n)!}|B_{2n}|t^{2n-1}\right)}{\sum_{n=1}^{\infty}\frac{2^{2n}-1}{(2n)!}2^{2n}p^{2n-1}|B_{2n}|t^{2n-1}}\\ &= \frac{\sum_{n=1}^{\infty}\frac{2^{2n}}{(2n)!}2n|B_{2n}|t^{2n-1}}{\sum_{n=1}^{\infty}\frac{2^{2n}-1}{(2n)!}2^{2n}p^{2n}|B_{2n}|t^{2n-1}}:=\frac{\sum_{n=1}^{\infty}a_nt^{2n-1}}{\sum_{n=1}^{\infty}b_nt^{2n-1}}\end{aligned}$$

where

$$a_n=\frac{2^{2n}}{(2n)!}2n|B_{2n}|,\ \ b_n=\frac{2^{2n}-1}{(2n)!}2^{2n}p^{2n}|B_{2n}|.$$

Clearly, if the monotonicity of a_n/b_n is proved, then by Lemma 3 it is deduced the monotonicity of f_1'/f_2', and then the monotonicity of the function F_p easily follows from Lemma 2. For this purpose, since a_n, $b_n>0$ for $n\in\mathbb{N}$, we only need to show that b_n/a_n is decreasing if $0<p\le1/\sqrt{5}$ and increasing if $1/2\le p\le1$. Indeed, elementary computation yields

$$\begin{aligned}\frac{b_{n+1}}{a_{n+1}}-\frac{b_n}{a_n} &= \frac{1}{2n+2}p^{2n+2}\left(2^{2n+2}-1\right)-\frac{1}{2n}p^{2n}\left(2^{2n}-1\right)\\ &= \frac{4^{n+1}-1}{2n+2}p^{2n}\left(p^2-\frac{n+1}{n}\frac{4^n-1}{4^{n+1}-1}\right).\end{aligned}$$

It is easy to obtain that for $n \in \mathbb{N}$

$$\frac{b_{n+1}}{a_{n+1}} - \frac{b_n}{a_n} \begin{cases} \geq 0 & \text{if } p^2 > \max_{n\in\mathbb{N}} \left(\frac{n+1}{n} \frac{4^n - 1}{4^{n+1} - 1} \right) = \frac{2}{5}, \\ \leq 0 & \text{if } p^2 \leq \min_{n\in\mathbb{N}} \left(\frac{n+1}{n} \frac{4^n - 1}{4^{n+1} - 1} \right) = \frac{1}{4}, \end{cases}$$

which proves the monotonicity of a_n/b_n.

By the monotonicity of the function F_p and the facts that

$$F_p\left(0^+\right) = \frac{2}{3p^2} \quad \text{and} \quad F_p\left(\frac{\pi}{2}^-\right) = -\frac{1}{\ln\left(\cos\left(\pi p/2\right)\right)},$$

the inequalities (2.5) and its reverse follow immediately. □

Lemma 6. *For every $t \in (0, \pi/2)$, $p \in (0, 1]$, the function G_p defined by*

$$G_p(t) = \frac{\ln \frac{\sin t}{t} + t \cot t - 1}{\ln \cos pt}$$

is increasing if $p \in (0, 1/2]$ and decreasing if $p \in [1/\sqrt{3}, 1]$. Consequently, for $p \in (0, 1/2]$ we have

$$\frac{1}{p^2} < \frac{\ln \frac{\sinh t}{t} + t \cot t - 1}{\ln \cos pt} < \frac{\ln 2 - \ln \pi - 1}{\ln\left(\cos\left(\pi p/2\right)\right)}. \tag{2.6}$$

It is reversed if $p \in [1/\sqrt{3}, 1]$.

Proof. We define $g_1(t) = \ln \frac{\sin t}{t} + t \cot t - 1$ and $g_2(t) = \ln(\cos pt)$, where $p \in (0, 1]$. Note that $g_1(0^+) = g_2(0^+) = 0$, then $G_p(t)$ can be written as

$$G_p(t) = \frac{g_1(t) - g_1(0^+)}{g_2(t) - g_2(0^+)}.$$

Differentiation and using (2.1) and (2.2) yield

$$\begin{aligned} \frac{g_1'(t)}{g_2'(t)} &= \frac{2\frac{\cos t}{\sin t} - \frac{1}{t} - t\frac{1}{\sin^2 t}}{-p \tan t} \\ &= \frac{2\left(\frac{1}{t} - \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} |B_{2n}| t^{2n-1}\right) - \frac{1}{t} - t\left(\frac{1}{t^2} + \sum_{n=1}^{\infty} \frac{(2n-1)2^{2n}}{(2n)!} |B_{2n}| t^{2n-2}\right)}{-p \sum_{n=1}^{\infty} \frac{2^{2n}-1}{(2n)!} 2^{2n} |B_{2n}| p^{2n-1} t^{2n-1}} \\ &= \frac{\sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} (2n+1) |B_{2n}| t^{2n-1}}{\sum_{n=1}^{\infty} \frac{2^{2n}-1}{(2n)!} 2^{2n} p^{2n} |B_{2n}| t^{2n-1}} := \frac{\sum_{n=1}^{\infty} c_n t^{2n-1}}{\sum_{n=1}^{\infty} d_n t^{2n-1}} \end{aligned}$$

where

$$c_n = \frac{2^{2n}}{(2n)!} (2n+1) |B_{2n}|, \quad d_n = \frac{2^{2n}-1}{(2n)!} 2^{2n} p^{2n} |B_{2n}|.$$

Similarly, we only need to show that d_n/c_n is decreasing if $0 < p \leq 1/\sqrt{5}$ and increasing if $1/2 \leq p \leq 1$. Indeed, elementary computation yields

$$\begin{aligned} \frac{d_{n+1}}{c_{n+1}} - \frac{d_n}{c_n} &= p^{2n+2} \frac{4^{n+1} - 1}{2n+3} - p^{2n} \frac{4^n - 1}{2n+1} \\ &= \frac{4^{n+1} - 1}{2n+3} p^{2n} \left(p^2 - \frac{2n+3}{2n+1} \frac{4^n - 1}{4^{n+1} - 1}\right). \end{aligned}$$

It is easy to obtain that for $n \in \mathbb{N}$

$$\frac{d_{n+1}}{c_{n+1}} - \frac{d_n}{c_n} \begin{cases} \geq 0 & \text{if } p^2 \geq \max_{n\in\mathbb{N}} \left(\frac{2n+3}{2n+1} \frac{4^n-1}{4^{n+1}-1} \right) = \frac{1}{3}, \\ \leq 0 & \text{if } p^2 \leq \min_{n\in\mathbb{N}} \left(\frac{2n+3}{2n+1} \frac{4^n-1}{4^{n+1}-1} \right) = \frac{1}{4}, \end{cases}$$

which proves the monotonicity of c_n/d_n.

By the monotonicity of the function F_p and the facts that

$$F_p\left(0^+\right) = \frac{1}{p^2} \quad \text{and} \quad F_p\left(\frac{\pi}{2}^-\right) = \frac{\ln 2 - \ln \pi - 1}{\ln\left(\cos\left(\pi p/2\right)\right)},$$

the inequalities (2.6) and its reverse follow immediately. □

Lemma 7 ([31], [13])**.** *For $t \in [0, \pi/2]$ and $p \in [0, 1)$, let $U_p(t)$ and $V_p(t)$ be defined by*

$$\begin{aligned} U_p(t) &= (\cos pt)^{1/p} \text{ if } p \neq 0 \text{ and } U_0(t) = 1, \\ V_p(t) &= (\cos pt)^{1/p^2} \text{ if } p \neq 0 \text{ and } V_0(t) = e^{-t^2/2}. \end{aligned}$$

Then both $U_p(t)$ and $V_p(t)$ are decreasing with p on $[0, 1)$.

3. Main results

Theorem 1. *For $t \in (0, \pi/2)$, the two-side inequality*

$$\left(\cos\frac{2t}{3}\right)^{3/2} < e^{t\cot t - 1} < (\cos p_1 t)^{1/p_1} \tag{3.1}$$

with the best constants $2/3$ and $p_1 \approx 0.6505$, where p_1 is the unique root of equation

$$1 + \frac{1}{p}\ln\left(\cos\frac{p\pi}{2}\right) = 0 \tag{3.2}$$

on $(0, 1)$. Moreover we have

$$\left(\cos\frac{2t}{3}\right)^{3/2} < e^{t\cot t - 1} < \left(\cos\frac{2t}{3}\right)^{1/\ln 2} < \frac{2\sqrt{2}}{e}\left(\cos\frac{2t}{3}\right)^{3/2}, \tag{3.3}$$

$$(\cos p_1 t)^{1/\left(3p_1^2\right)} < e^{t\cot t - 1} < (\cos p_1 t)^{1/p_1}, \tag{3.4}$$

where the exponents $3/2$ and $1/\ln 2$ in (3.3) are the best constants, and $p_1 \approx 0.6505$ in (3.4) is also the best.

Proof. (i) We first prove the first inequality in (3.1) holds for $t \in (0, \pi/2)$ and $2/3$ is the best constant. Letting $p = 2/3 \in [\sqrt{10}/5, 1]$ in (2.5) yields the first one in (3.1) and the second one in (3.3). If there is a $p < 2/3$ such that $e^{t\cot t - 1} > (\cos pt)^{1/p}$ holds for $t \in (0, \pi/2)$, then we have

$$\lim_{t\to 0^+} \frac{t\cot t - 1 - \frac{1}{p}\ln(\cos pt)}{t^2} > 0.$$

Using power series extension gives

$$t\cot t - 1 - \frac{1}{p}\ln(\cos pt) = t^2\left(\frac{1}{2}p - \frac{1}{3}\right) + O\left(t^4\right),$$

and therefore,

$$\lim_{t\to 0^+} \frac{t\cot t - 1 - \frac{1}{p}\ln(\cos pt)}{t^2} = \frac{1}{2}\left(p - \frac{2}{3}\right) > 0,$$

which yields a contradiction. Hence, $2/3$ is the best possible.

(ii) By Lemma 7, it is seen that the function $p \mapsto 1+\frac{1}{p}\ln\left(\cos\frac{p\pi}{2}\right)$ is decreasing on $(0,1)$, which together with the facts that

$$\lim_{p\to 0^+}\left(1+\frac{1}{p}\ln\left(\cos\frac{p\pi}{2}\right)\right)=1 \quad\text{and}\quad \lim_{p\to 1^-}\left(1+\frac{1}{p}\ln\left(\cos\frac{p\pi}{2}\right)\right)=-\infty$$

reveals that the equation (3.2) has a unique root $p_1\in(0,1)$. Numerical calculation gives $p_1\approx 0.6505$. Letting $p=p_1\in[\sqrt{10}/5,1]$ in Lemma 5 yields

$$-\frac{1}{\ln\left(\cos\left(\pi p_1/2\right)\right)}<\frac{t\cot t-1}{\ln\left(\cos p_1 t\right)}<\frac{2}{3p_1^2}.$$

The first inequality above can be reduced as

$$\frac{2}{3p_1^2}\ln\left(\cos p_1 t\right)<t\cot t-1<-\frac{1}{\ln\left(\cos\left(\pi p_1/2\right)\right)}\ln\left(\cos p_1 t\right)=\frac{1}{p_1}\ln\left(\cos p_1 t\right),$$

that is, the second one in (3.1) and the first one in (3.4), where the equality is due to that p_1 is the unique root of equation (3.2). It is clear that p_1 is the best.

(iii) It remains to prove the third one in (3.3). To this end, it suffices to check that

$$\frac{1}{\ln 2}\ln\left(\cos\frac{2t}{3}\right)-\ln\frac{2\sqrt{2}}{e}-\frac{3}{2}\ln\left(\cos\frac{2t}{3}\right)<0.$$

Since $\ln\left(\cos\frac{2t}{3}\right)>\ln\left(\cos\frac{\pi}{3}\right)=\ln 2$, the left-hand side is less than

$$\left(\frac{1}{\ln 2}-\frac{3}{2}\right)\ln 2-\ln\frac{2\sqrt{2}}{e}=2-3\ln 2<0,$$

which proves the desired inequality.

This completes the proof. □

Yang has shown in [32] that the inequalities

$$\left(\cos pt\right)^{1/p}<\frac{\sin t}{t}<\left(\cos qt\right)^{1/q}$$

hold for $t\in(0,\pi/2)$ if and only if $p\in[p_0,1)$ and $q\in(0,1/3]$, where $p_0\approx 0.3473$. Hence, we have

$$\cos t<\left(\cos\frac{t}{2}\right)^2<\left(\cos p_0 t\right)^{1/p_0}<\frac{\sin t}{t}<\left(\cos\tfrac{t}{3}\right)^3<\frac{2+\cos t}{3}. \tag{3.5}$$

On the other hand, utilizing Theorem 1 and Lemma 7 we derive

$$\cos t<\left(\cos\frac{3t}{4}\right)^{4/3}<\left(\cos\frac{2t}{3}\right)^{3/2}<e^{t\cot t-1}<\left(\cos p_1 t\right)^{1/p_1}<\left(\cos\frac{t}{2}\right)^2<\left(\cos\frac{t}{3}\right)^3<1,$$

which in combination with (3.5) leads to the following

Corollary 1. *For $t\in(0,\pi/2)$, the chain of inequalities hold:*

(3.6)

$$\cos t<\left(\cos\tfrac{3t}{4}\right)^{4/3}<\left(\cos\tfrac{2t}{3}\right)^{3/2}<e^{t\cot t-1}<\left(\cos\tfrac{t}{2}\right)^2<\frac{\sin t}{t}<\left(\cos\tfrac{t}{3}\right)^3<\tfrac{2+\cos t}{3}.$$

By Lemma 5 and Lemma 7 we also have

Theorem 2. *For $t \in (0, \pi/2)$, the two-side inequality*

$$(\cos pt)^{2/\left(3p^2\right)} < e^{t\cot t - 1} < (\cos qt)^{2/\left(3q^2\right)} \tag{3.7}$$

holds if $p \in [\sqrt{10}/5, 1]$ and $q \in (0, 1/2]$.

In particular, letting $p = 2/3, q = 1/2, 1/3, 0^+$, we have

$$\left(\cos\frac{2t}{3}\right)^{3/2} < e^{t\cot t - 1} < \left(\cos\frac{t}{2}\right)^{8/3} < \left(\cos\frac{t}{3}\right)^{6} < e^{-t^2/3}. \tag{3.8}$$

Taking $p = 1/2$ in (2.5) leads to

$$\left(\cos\frac{t}{2}\right)^{2/\ln 2} < e^{t\cot t - 1} < \left(\cos\frac{t}{2}\right)^{8/3}, \tag{3.9}$$

where the exponents $2/\ln 2$ and $8/3$ are the best possible.

Replacing t, p, q by $t/2$, $2p$, $2q$ in Theorem 2 and next taking squares, we get

Corollary 2. *For $t \in (0, \pi)$, the two-side inequality*

$$(\cos pt)^{1/\left(3p^2\right)} < e^{t\cot(t/2) - 2} < (\cos qt)^{1/\left(3q^2\right)} \tag{3.10}$$

holds if $p \in [1/\sqrt{10}, 1/2]$ and $q \in (0, 1/4]$. Moreover, we have

$$\left(\cos\frac{t}{3}\right)^{1/3} < e^{t\cot(t/2) - 2} < \left(\cos\frac{t}{4}\right)^{16/3} < \left(\cos\frac{t}{6}\right)^{12} < e^{-t^2/6}, \tag{3.11}$$

$$\left(\cos\frac{t}{4}\right)^{4/\ln 2} < e^{t\cot(t/2) - 2} < \left(\cos\frac{t}{4}\right)^{16/3}. \tag{3.12}$$

Theorem 3. *For $t \in (0, \pi/2)$, we have*

$$\cos^{2/3} t < \frac{2}{3}\cos t + \frac{1}{3} < \left(\cos\frac{2t}{3}\right)^{3/2}, \tag{3.13}$$

$$\left(\cos\frac{t}{2}\right)^{4/3} < \frac{\sin t}{t} < \frac{2\cos\frac{t}{2} + \cos^2\frac{t}{2}}{3} < \left(\frac{2}{3}\cos\frac{t}{2} + \frac{1}{3}\right)^2 < \cos^3\frac{t}{3}. \tag{3.14}$$

Proof. (i) The firs inequality in (3.13) is from the basic inequality. To prove the second one in (3.13), we define

$$f(t) = \left(\frac{2}{3}\cos t + \frac{1}{3}\right)\left(\cos\frac{2t}{3}\right)^{-3/2}.$$

Differentiation and simplifying leads to

$$f'(t) = -\frac{2}{3}\left(\cos\frac{2t}{3}\right)^{-5/2}\left(\sin\frac{t}{3}\right)\left(1 - \cos\frac{t}{3}\right) < 0.$$

Therefore we have $f(t) < f(0) = 1$, which proves (3.13).

(ii) Now we prove (3.14). The first and second ones are due to Neuman and Sándor [17, (2.5)]. The third one easily follows from

$$\frac{2\cos\frac{t}{2} + \cos^2\frac{t}{2}}{3} - \left(\frac{2}{3}\cos\frac{t}{2} + \frac{1}{3}\right)^2 = -\frac{1}{9}\left(\cos\frac{t}{2} - 1\right)^2 < 0.$$

Replacing $2t$ by t and taking square values in the second one of (3.13) yields that the fourth one holds for $t \in (0, \pi)$.

Thus the proof is complete. □

Taking square roots for each sides in (3.13) and the first and second inequalities of (3.8), we get

$$(\cos t)^{1/3} < \sqrt{\tfrac{2}{3}\cos t + \tfrac{1}{3}} < \left(\cos \tfrac{2t}{3}\right)^{3/4} < \sqrt{e^{t\cot t - 1}} < \left(\cos \tfrac{t}{2}\right)^{4/3}.$$

This in combination with (3.14), (3.11) and

$$e^{-t^2/6} < \frac{2+\cos t}{3} \quad \text{for } t \in (0,\infty)$$

proved in [31, Theorem 2], [36] leads to the following interesting chain of inequalities for trigonometric functions.

Corollary 3. *For $t \in (0, \pi/2)$, we have*

$$\begin{aligned}(\cos t)^{1/3} &< \sqrt{\tfrac{2}{3}\cos t + \tfrac{1}{3}} < \left(\cos \tfrac{2t}{3}\right)^{3/4} < \sqrt{e^{t\cot t - 1}} < \\ \left(\cos \tfrac{t}{2}\right)^{4/3} &< \frac{\sin t}{t} < \frac{2\cos\frac{t}{2} + \cos^2\frac{t}{2}}{3} < \left(\tfrac{2}{3}\cos\tfrac{t}{2} + \tfrac{1}{3}\right)^2 < \\ \left(\cos\tfrac{t}{3}\right)^3 &< e^{t\cot\frac{t}{2} - 2} < \left(\cos\tfrac{t}{4}\right)^{16/3} < \left(\cos\tfrac{t}{6}\right)^{12} < e^{-t^2/6} < \tfrac{2}{3} + \tfrac{1}{3}\cos t.\end{aligned} \tag{3.15}$$

The fourth and fifth inequalities in (3.15) implies that for $t \in (0, \pi/2)$,

$$e^{t\cot t - 1} < \left(\cos\tfrac{t}{2}\right)^{8/3} < \left(\frac{\sin t}{t}\right)^2. \tag{3.16}$$

Further, we have

Corollary 4. *For $t \in (0, \pi/2)$, we have*

$$\begin{aligned}\frac{\pi^2}{4e}\left(\frac{\sin t}{t}\right)^2 &< \left(\frac{\sin t}{t}\right)^{1/(\ln\pi - \ln 2)} < \left(\cos\frac{t}{2}\right)^{2/\ln 2} < e^{t\cot t - 1} \\ &< \left(\cos\tfrac{t}{2}\right)^{8/3} < \left(\frac{\sin t}{t}\right)^2 < \frac{2^{10/3}}{\pi^2}\left(\cos\tfrac{t}{2}\right)^{8/3},\end{aligned} \tag{3.17}$$

$$\frac{\sin t}{t} < \tfrac{2}{3} + \tfrac{1}{3}\cos t < \frac{e^{t\cot t - 1} + 1}{2}, \tag{3.18}$$

$$e^{t\cot t - 1} > \tfrac{2}{3}\cos t + \tfrac{1}{3} > \frac{\frac{\sin t}{t} + \cos t}{2}. \tag{3.19}$$

Proof. (i) We first prove (3.17). To obtain the first inequality in (3.17), it is enough to check that

$$\frac{1}{\ln\pi - \ln 2}\ln\frac{\sin t}{t} - \left(\ln\frac{\pi^2}{4e} + 2\ln\frac{\sin t}{t}\right) > 0.$$

Using Jordan inequality $(\sin t)/t > 2/\pi$ for $t \in (0, \pi/2)$ we see that the left-hand side above is greater than

$$\left(\frac{1}{\ln\pi - \ln 2} - 2\right)\ln\frac{2}{\pi} - \ln\frac{\pi^2}{4e} = 0,$$

which proves the first one.

Using the known inequalities that for $x \in (0, \pi/2)$

$$\left(\cos\frac{x}{2}\right)^{4/3} < \frac{\sin x}{x} < \left(\cos\frac{x}{2}\right)^{2(\ln\pi - \ln 2)/\ln 2} \tag{3.20}$$

proved in [15] by Lv et al. gives the second and fifth ones.

The third and fourth ones are (3.12). While the last one is the right-hand side one in [31, (3.15)], [36].

(ii) Now we prove ((3.18)). The first inequality in (3.18) has proven previously, while the second one is obtained from the last two ones in (3.6), that is, the well-known Cusa's inequality. The third one is equivalent to

$$\tfrac{2}{3}\cos t+\tfrac{1}{3}<e^{t\cot t-1}, \tag{3.21}$$

which easily follows from the second and third ones in (3.15).

(iii) We show (3.19) at last. The first inequality is (3.21), while the second one is equivalent to the second one in (3.18), which proves (3.19) and the proof is completed. □

By Lemma 6 we have

Theorem 4. *For $t\in(0,\pi/2)$, the two-side inequality*

$$(\cos pt)^{1/(2p^2)}<\sqrt{\frac{\sin t}{t}\exp\left(t\cot t-1\right)}<(\cos qt)^{1/(2q^2)} \tag{3.22}$$

holds if $p\in[1/\sqrt{3},1]$ and $q\in(0,1/2]$. Moreover, we have

$$\sqrt{\frac{8}{\pi e}}\cos^2\frac{t}{2}\ <\ \cos^{\beta}\frac{t}{2}<\sqrt{\frac{\sin t}{t}\exp\left(t\cot t-1\right)}<\cos^2\frac{t}{2}, \tag{3.23}$$

$$\cos^{3/2}\frac{t}{\sqrt{3}}\ <\ \sqrt{\frac{\sin t}{t}\exp\left(t\cot t-1\right)}<\cos^{\gamma}\frac{t}{\sqrt{3}}, \tag{3.24}$$

$$\sqrt{\cos t}\ <\ \sqrt{\frac{\sin t}{t}\exp\left(t\cot t-1\right)}<\cos^2\frac{t}{2} \tag{3.25}$$

where the exponents

$$\beta=\frac{\ln\pi-\ln 2+1}{\ln 2}\approx 2.0942,\ 2\ \ \textit{and}\ \ 3/2,\ \gamma=-\frac{(\ln\pi-\ln 2+1)}{2\ln(\cos\frac{\pi}{2\sqrt{3}})}\approx 1.4990$$

are the best constants.

Proof. (i) The first result in this theorem is a direct consequence of Lemma 6.

(ii) Letting $p=1/2\in(0,1/2]$ in (2.6) yields the second and third inequalities in (3.23). To show the first one, it suffices to check that

$$\frac{1}{2}\ln\frac{8}{\pi e}+2\ln\cos\frac{t}{2}-\frac{\ln\pi-\ln 2+1}{\ln 2}\ln\cos\frac{t}{2}<0.$$

Since $\ln\cos\frac{t}{2}>\ln\cos\frac{\pi}{4}=-\frac{1}{2}\ln 2$, the left-hand side is less than

$$\frac{1}{2}\ln\frac{8}{\pi e}-\left(2-\frac{\ln\pi-\ln 2+1}{\ln 2}\right)\frac{1}{2}\ln 2=0,$$

which proves (3.23).

(iii) Letting $p=1/\sqrt{3}\in[1/\sqrt{3},1]$ in (2.6) yields (3.24).

(iv) Taking $p=1$ and $q=1/2$ in (3.22) gives (3.25).

This proves the desired results. □

Remark 1. *From the second inequality in (3.25) in conjunction with (3.16) it is seen that*

$$\sqrt{e^{t\cot t-1}}<\left(\frac{\sin t}{t}e^{t\cot t-1}\right)^{1/3}<\left(\cos\tfrac{t}{2}\right)^{4/3}<\frac{\sin t}{t}.$$

Theorem 5. *For $t \in (0, \pi/2)$, we have*

(3.26)

$$\frac{\frac{e(\pi-2)}{\pi}e^{t\cot t-1}+\frac{\sin t}{t}}{2}<\frac{1+\cos t}{2}<\frac{\frac{\sin t}{t}+e^{t\cot t-1}}{2}<\left(e^{-1}+\tfrac{2}{\pi}\right)\frac{1+\cos t}{2}.$$

Proof. (i) We first prove the first and second inequalities in (3.26). For this purpose, let us define

$$g(t)=(t\cot t-1)-\ln\left(1+\cos t-\frac{\sin t}{t}\right).$$

Differentiating $g(t)$ gives

$$g'(t)=\frac{t-\sin t}{t(t-\sin t+t\cos t)}g_1(t),$$

where

$$g_1(t)=-\frac{\cos t+1}{\sin^2 t}t^2+\frac{t}{\sin t}+1.$$

Using double angle formula and Lemma 4 we have

$$\begin{aligned} g_1(t) &= -\frac{t^2}{2\sin^2\frac{t}{2}}+\frac{t}{\sin t}+1 \\ &= -\tfrac{t^2}{2}\left(\tfrac{1}{\left(\frac{t}{2}\right)^2}+\sum_{n=1}^{\infty}\tfrac{(2n-1)2^{2n}}{(2n)!}|B_{2n}|\left(\tfrac{t}{2}\right)^{2n-2}\right)+t\left(\tfrac{1}{t}+\sum_{n=1}^{\infty}\tfrac{2^{2n}-2}{(2n)!}|B_{2n}|t^{2n-1}\right)+1 \\ &= \sum_{n=1}^{\infty}\frac{4^{n-1}-n}{(2n)!}|B_{2n}|t^{2n}>0. \end{aligned}$$

Hence, $g'(t)>0$ for $t\in(0,\pi/2)$, and so

$$0=\lim_{t\to0^+}g(t)<g(t)<\lim_{t\to\pi/2^-}g(t)=\ln\frac{\pi}{e(\pi-2)},$$

which implies the desired inequalities.

(ii) Now we prove the second and third ones by defining

$$h(t)=\frac{\frac{\sin t}{t}+e^{t\cot t-1}}{1+\cos t}.$$

Differentiation leads to

$$\begin{aligned} h'(t) &= \frac{-(\cos t+1)(t-\sin t)}{\left(\sin^2 t\right)(\cos t+1)^2}e^{t\cot t-1}+\frac{(\cos t+1)(t-\sin t)}{t^2(\cos t+1)^2} \\ &= \frac{t-\sin t}{\left(\sin^2 t\right)(\cos t+1)^2}\left(\frac{\sin^2 t}{t^2}-e^{t\cot t-1}\right). \end{aligned}$$

It is obtained by 3.16 that $h'(t)>0$ for $t\in(0,\pi/2)$, and therefore, we conclude that

$$1=\lim_{t\to0^+}h(t)<h(t)<\lim_{t\to\pi/2^-}h(t)=e^{-1}+\frac{2}{\pi},$$

which deduces the second and third ones in ((3.26). □

Remark 2. *In the same way as part two of Theorem 5, we can prove a new two-side inequality for hyperbolic functions:*

$$\frac{1+\cosh t}{e}<\frac{\frac{\sinh t}{t}+e^{\frac{t\cosh t}{\sinh t}-1}}{2}<\frac{1+\cosh t}{2}.$$

In fact, we define

$$k\left(t\right)=\frac{\frac{\sinh t}{t}+e^{\frac{t\cosh t}{\sinh t}-1}}{1+\cosh t}.$$

Differentiation and simplifying give us

$$k'\left(t\right)=\frac{\sinh t-t}{\left(\cosh t+1\right)\sinh^{2}t}\left(e^{t\frac{\cosh t}{\sinh t}-1}-\frac{\sinh^{2}t}{t^{2}}\right)<0,$$

where the inequality holds is due to the first one in (1.3) and $\sinh t>t$ *for* $t>0$. *It follows that*

$$2e^{-1}=\lim_{t\to\infty}k\left(t\right)<k\left(t\right)<\lim_{t\to0}k\left(t\right)=1,$$

which is desired inequality. Clearly, this method is simpler than Alzer's, and our result is also more nice.

Employing inequalities (3.25) and (3.26), we immediately the trigonometric version of (1.5).

Corollary 5. *For* $t\in(0,\pi/2)$, *we have*

$$\sqrt{\cos t}<\left(\frac{\sin t}{t}e^{t\cot t-1}\right)^{1/2}<\frac{1+\cos t}{2}<\frac{\frac{\sin t}{t}+e^{t\cot t-1}}{2}<\left(e^{-1}+\frac{2}{\pi}\right)\frac{1+\cos t}{2},$$

Lastly, we give the trigonometric versions of (1.6) and (1.7).

Theorem 6. *Let* $t\in(0,\pi/2)$. *Then*

(i) if $p\geq6/5$, *then the two-side inequality*

$$\alpha\cos^{p}t+(1-\alpha)<e^{p(t\cot t-1)}<\beta\cos^{p}t+(1-\beta)\tag{3.27}$$

holds if and only if $\alpha\geq(1-e^{-p})$ *and* $\beta\leq2/3$*;*

(ii) if $0<p\leq1$, *then (3.27) holds if and only if* $\alpha\geq2/3$ *and* $\beta\leq(1-e^{-p})$*;*

(iii) if $p<0$, *then (3.27) holds if and only if* $\alpha\leq0$ *and* $\beta\geq2/3$*;*

(iv) the double inequality

$$M_{p}\left(\cos t,1;\tfrac{2}{3}\right)<e^{t\cot t-1}<M_{q}\left(\cos t,\tfrac{2}{3}\right)\tag{3.28}$$

holds if and only if $p\leq\ln3$ *and* $q\geq6/5$, *where* $M_{p}\left(x,y;w\right)$ $(w\in(0,1))$ *is the weighted power mean of order* r *of positive* x *and* y *defined by*

$$M_{r}\left(x,y;w\right)=\left(wx^{r}+\left(1-w\right)y^{r}\right)^{1/r}\ \text{ if } r\neq0 \text{ and } M_{0}\left(x,y;w\right)=x^{w}y^{1-w}\tag{3.29}$$

Proof. For $t\in(0,\pi/2)$ and $p\neq0$, we define

$$u\left(t\right)=\frac{1-e^{p(t\cot t-1)}}{1-\cos^{p}t}:=\frac{u_{1}\left(t\right)}{u_{2}\left(t\right)}.$$

Since $u_{1}\left(0^{+}\right)=u_{2}\left(0^{+}\right)=0$, $u\left(t\right)$ can be written as

$$u\left(t\right)=\frac{u_{1}\left(t\right)-u_{1}\left(0^{+}\right)}{u_{2}\left(t\right)-u_{2}\left(0^{+}\right)}.\tag{3.30}$$

Differentiation gives

$$\frac{u_{1}'\left(t\right)}{u_{2}'\left(t\right)}=\frac{e^{p(t\cot t-1)}\frac{t-\cos t\sin t}{\sin^{2}t}}{\cos^{p-1}t\sin t}:=u_{3}\left(t\right),\tag{3.31}$$

$$u_{3}'\left(t\right)=\frac{e^{p(t\cot t-1)}}{\sin^{5}t\cos^{p}t}\left(p\times u_{4}\left(t\right)-u_{5}\left(t\right)\right)=\frac{e^{p(t\cot t-1)}}{\sin^{5}t\cos^{p}t}u_{4}\left(t\right)\left(p-\frac{u_{5}\left(t\right)}{u_{4}\left(t\right)}\right),\tag{3.32}$$

where

$$\begin{aligned} u_4(t) &= -t^2\cos t\sin^2 t - t^2\cos^3 t + t\cos^2 t\sin t + t\cos^2 t\sin t + t\sin^3 t - \cos t\sin^2 t, \\ u_5(t) &= t\left(3\cos^2 t\sin t + \sin^3 t\right) - 3\cos t\sin^2 t. \end{aligned}$$

Clearly, if we prove $u_3'(t) > 0$ if $p \geq 6/5$ and $u_3'(t) < 0$ if $p \leq 1$ with $p \neq 0$, then by Lemma 2 we see that u is increasing if $p \geq 6/5$ and decreasing if $p \leq 1$ with $p \neq 0$. And then, we can derive that

$$\begin{aligned} \tfrac{2}{3} &= \lim_{t\to 0^+} u(t) < u(t) = \frac{1-e^{p(t\cot t-1)}}{1-\cos^p t} < \lim_{t\to\pi/2^-} u(t) = 1-e^{-p} \text{ for } p \geq 6/5, \\ 1-e^{-p} &= \lim_{t\to\pi/2^-} u(t) < u(t) = \frac{1-e^{p(t\cot t-1)}}{1-\cos^p t} < \lim_{t\to 0^+} u(t) = \tfrac{2}{3} \text{ for } 0 < p \leq 1, \\ 0 &= \lim_{t\to\pi/2^-} u(t) < u(t) = \frac{1-e^{p(t\cot t-1)}}{1-\cos^p t} < \lim_{t\to 0^+} u(t) = \tfrac{2}{3} \text{ for } p < 0, \end{aligned}$$

which yield the first, second and third results in this theorem.

Now we will show that $u_3'(t) > 0$ if $p \geq 6/5$ and $u_3'(t) < 0$ if $p \leq 1$ with $p \neq 0$. Simplifying yields

$$u_4(t) = (\sin t - t\cos t)(t - \cos t\sin t) > 0$$

for $t \in (0, \pi/2)$. Using (2.2)–(2.4) and simplifying we have

$$\begin{aligned} \frac{u_5(t)}{\cos t\sin^2 t} &= 3t\frac{\cos t}{\sin t} + t\frac{\sin t}{\cos t} - 3 = \sum_{n=1}^{\infty} \frac{4^n-4}{(2n)!} 2^{2n}|B_{2n}|t^{2n}, \\ \frac{u_4(t)}{\cos t\sin^2 t} &= 2t\frac{\cos t}{\sin t} - t^2\frac{1}{\sin^2 t} + t\frac{\sin t}{\cos t} - 1 = \sum_{n=1}^{\infty} (4^n - 2n - 2)\frac{2^{2n}}{(2n)!}|B_{2n}|t^{2n}, \end{aligned}$$

and then, we get

$$\frac{u_5(t)}{u_4(t)} = \frac{\sum_{n=1}^{\infty} \frac{4^n-4}{(2n)!} 2^{2n}|B_{2n}|t^{2n}}{\sum_{n=1}^{\infty} (4^n-2n-2)\frac{2^{2n}}{(2n)!}|B_{2n}|t^{2n}} := \frac{\sum_{n=2}^{\infty} a_n t^{2n}}{\sum_{n=2}^{\infty} b_n t^{2n}}.$$

By Lemma 3, in order to prove the monotonicity of $u_5(t)/u_4(t)$, it suffices to determine the monotonicity of a_n/b_n. We have

$$\frac{a_n}{b_n} = \frac{4^n-4}{4^n-2n-2} := c(n).$$

Differentiating $c(x)$ we get

$$c'(x) = -2\frac{4^x((x-1)\ln 4 - 1) + 4}{(4^x-2x-2)^2} < 0$$

for $x \geq 2$. Then $t \mapsto u_5(t)/u_4(t)$ is decreasing on $(0, \pi/2)$, and we conclude that

$$1 = \lim_{t\to\pi/2^-} \frac{u_5(t)}{u_4(t)} < \frac{u_5(t)}{u_4(t)} < \lim_{t\to 0^+} \frac{u_5(t)}{u_4(t)} = \frac{6}{5}.$$

Thus, $u_3'(t) > 0$ if $p \geq 6/5$ and $u_3'(t) < 0$ if $p \leq 1$ with $p \neq 0$.

At last, we prove the fourth result. The first one implies that the right-hand side inequality in (3.28) holds if $q \geq 6/5$. While the necessity is obtained from the following limit relation:

$$\lim_{t\to 0^+} \frac{t\cot t - 1 - \ln M_q\left(\cos t, 1; \tfrac{2}{3}\right)}{t^4} \leq 0.$$

Expanding in power series leads to

$$t\cot t-1-\ln M_q\left(\cos t,1;\tfrac{2}{3}\right)=-\frac{1}{36}\left(q-\frac{6}{5}\right)t^4+O\left(t^6\right).$$

It is deduced that $q\geq 6/5$.

We now prove the left-hand side inequality holds if and only if $p\leq\ln 3$. The necessity easily follows from

$$\lim_{t\to\pi/2^-}\left(t\cot t-1-\ln M_p\left(\cos t,1;\tfrac{2}{3}\right)\right)=\begin{cases}-1+\frac{1}{p}\ln 3 & \text{if } p>0,\\ \infty & \text{if } p\leq 0\end{cases}\geq 0.$$

Next we deal with the sufficiency. We distinguish two cases.

Case 1: $p\leq 1$. By the second and third results proved previously, the sufficiency is immediate in this case.

Case 2: $p\in(1,\ln 3]$. As shown previously, the functions $t\mapsto u_5(t)/u_4(t)$ is decreasing on $(0,\pi/2)$, and so the function $t\mapsto(p-u_5(t)/u_4(t)):=u_6(t)$ is increasing on the same interval. This together with the facts that

$$u_6\left(0^+\right)=p-\frac{6}{5}<0\ \text{ and }\ u_6\left(\tfrac{\pi}{2}^-\right)=p-1>0$$

indicates that there is a unique $t_0\in(0,\pi/2)$ such that $u_6(t)<0$ for $t\in(0,t_0)$ and $u_6(t)>0$ for $t\in(t_0,\pi/2)$, which by (3.32) in turn reveals that u_3 is decreasing on $(0,t_0)$ and increasing on $(t_0,\pi/2)$. From Lemma 2 it is seen that u is decreasing on $(0,t_0)$, and so we have

$$u(t_0)\leq u(t)=\frac{1-e^{p(t\cot t-1)}}{1-\cos^p t}<u\left(0^+\right)=\frac{2}{3}\text{ for } t\in(0,t_0],$$

which can be changed into

$$e^{p(t\cot t-1)}>\frac{2}{3}\cos^p t+\frac{1}{3}\text{ for } t\in(0,t_0]. \tag{3.33}$$

On the other hand, Lemma 2 also implies that

$$t\mapsto\frac{u_1(t)-u_1\left(\frac{\pi}{2}^-\right)}{u_2(t)-u_2\left(\frac{\pi}{2}^-\right)}=\frac{e^{p(t\cot t-1)}-e^{-p}}{\cos^p t}:=v(t)$$

is increasing on $(t_0,\pi/2)$, and therefore, we get

$$v(t)=\frac{e^{p(t\cot t-1)}-e^{-p}}{\cos^p t}>\frac{e^{p(t_0\cot t_0-1)}-e^{-p}}{\cos^p t_0}\text{ for } t\in(t_0,\pi/2),$$

which implies that

$$e^{p(t\cot t-1)}>\frac{e^{p(t_0\cot t_0-1)}-e^{-p}}{\cos^p t_0}\cos^p t+e^{-p}\text{ for } t\in(t_0,\pi/2). \tag{3.34}$$

Clearly, if we prove the right-hand side in (3.34) is also greater than the one in (3.33), then the proof will be complete. Since t_0 satisfies (3.33), for $t\in(t_0,\pi/2)$, we have

$$\begin{aligned}e^{p(t\cot t-1)} &> >\frac{e^{p(t_0\cot t_0-1)}-e^{-p}}{\cos^p t_0}\cos^p t+e^{-p}>\frac{\left(\frac{2}{3}\cos^p t_0+\frac{1}{3}\right)-e^{-p}}{\cos^p t_0}\cos^p t+e^{-p}\\ &= \frac{2}{3}\cos^p t+\frac{1}{3}+\left(e^{-p}-\frac{1}{3}\right)\frac{\cos^p t_0-\cos^p t}{\cos^p t_0}\geq\frac{2}{3}\cos^p t+\frac{1}{3},\end{aligned}$$

where the last inequality holds is due to $p\in(1,\ln 3]$ and $t\in(t_0,\pi/2)$.

Thus the proof is finished. □

4. The corresponding inequalities for means

For $a, b > 0$ with $a \neq b$, let L, Q, A, G, P and T stand for the logarithmic,, quadratic,, arithmetic, geometric, the first and second means [23], [24] of a and b defined by

$$Q = Q(a,b) = \sqrt{\tfrac{a^2+b^2}{2}}, \quad A = A(a,b) = \tfrac{a+b}{2}, \quad G = G(a,b) = \sqrt{ab},$$

$$P = P(a,b) = \frac{a-b}{2\arcsin\frac{a-b}{a+b}}, \quad T = T(a,b) = \frac{a-b}{2\arctan\frac{a-b}{a+b}},$$

respectively. Very recently, Sándor present a new mean in [22] defined as

$$X = X(a,b) = Ae^{G/P-1}, \tag{4.1}$$

where A, G, P are the arithmetic, geometric and the first Seiffert means as defined previously. In what follows, we also can encounter other three functions defined by

$$B = B(a,b) = Qe^{A/T-1} \tag{4.2}$$

$$J = J(a,b) = Ae^{(A+G)/P-2}, \tag{4.3}$$

$$K = K(a,b) = Qe^{(Q+A)/T-2}, \tag{4.4}$$

where Q, A, T are the quadratic, arithmetic and the second Seiffert means, respectively. They will be showed to be means of a and b in the sequel.

There has many inequalities for these means, we quote [16], [18], [8], [7], [9], [10], [11], [29], [33], [34], [35].

In order to obtain inequalities for these means corresponding to ones established in previous section, we need two variable substitutions:
(i) Substitution 1: Letting $t = \arcsin\frac{b-a}{a+b}$ yields

$$\frac{\sin t}{t} = \frac{P}{A}, \cos t = \frac{G}{A}, e^{t\cot t-1} = e^{G/P-1}, e^{t\cot(t/2)-2} = e^{(A+G)/P-2}; \tag{4.5}$$

(ii) Substitution 2: Letting $t = \arctan\frac{b-a}{a+b}$ yields

$$\frac{\sin t}{t} = \frac{T}{Q}, \cos t = \frac{A}{Q}, e^{t\cot t-1} = e^{A/T-1}, e^{t\cot(t/2)-2} = e^{(Q+A)/T-2}. \tag{4.6}$$

For simplicity in expressions, however, we only select those functions composed by $(\sin t)/t$, $\cos t$, $\cos(t/2)$, $\cos(t/4)$ in a chain of inequalities given in previous section. For example, from Corollary 1 we get

$$\cos t < e^{t\cot t-1} < \frac{1+\cos t}{2} < \frac{\sin t}{t} < \frac{2+\cos t}{3}.$$

Using Substitution 1 and next multiplying all functions by A yield

$$G < A\exp\left(\frac{G}{P}-1\right) = X < \frac{A+G}{2} < P < \frac{2A+G}{3}. \tag{4.7}$$

Using Substitution 2 and next multiplying all functions by Q yield

$$A < Q\exp\left(\frac{A}{T}-1\right) = B < \frac{Q+A}{2} < T < \frac{2Q+A}{3}. \tag{4.8}$$

The above two inequalities can be stated as a proposition.

Proposition 1. *For $a, b > 0$ with $a \neq b$, both the inequalities (4.7) and (4.8) are true.*

Remark 3. *From (4.8) it is easy to find that the function B defined by (4.2) is indeed a mean of a and b, and is between A and Q.*

Inequalities (3.9) can be changed into

Proposition 2. *For $a, b > 0$ with $a \neq b$, both the two-side inequalities*

$$\left(\frac{A+G}{2}\right)^{1/\ln 2} A^{1-1/\ln 2} \quad < \quad X < \left(\frac{A+G}{2}\right)^{4/3} A^{-1/3}, \tag{4.9}$$

$$\left(\frac{Q+A}{2}\right)^{1/\ln 2} Q^{1-1/\ln 2} \quad < \quad B < \left(\frac{A+G}{2}\right)^{4/3} A^{-1/3} \tag{4.10}$$

hold.

Applying Substitutions 1 and 2 to (3.12) we have

Proposition 3. *For $a, b > 0$ with $a \neq b$, both the two-side inequalities*

$$\left(\tfrac{\sqrt{2A}+\sqrt{A+G}}{2\sqrt{2}}\right)^{2/\ln 2} A^{1-1/\ln 2} \quad < \quad J < \left(\tfrac{\sqrt{2A}+\sqrt{A+G}}{2\sqrt{2}}\right)^{8/3} A^{-1/3}, \tag{4.11}$$

$$\left(\tfrac{\sqrt{2Q}+\sqrt{Q+A}}{2\sqrt{2}}\right)^{2/\ln 2} Q^{1-1/\ln 2} \quad < \quad K < \left(\tfrac{\sqrt{2Q}+\sqrt{Q+A}}{2\sqrt{2}}\right)^{8/3} A^{-1/3} \tag{4.12}$$

are true.

Remark 4. *From (4.11) and (4.12) it is easy to see that the function J, K defined by (4.3), 4.4 are indeed means of a and b, and satisfy the relations: $G < J < A$ and $A < K < Q$.*

In the sequel, we only list a chain of inequalities for means deduced from Substitution 1, since another one can be clearly derived via replacing (G, A, P, X, J) by (A, Q, T, B, K).

The chain of inequalities (3.15) can be changed into

Proposition 4. *For $a, b > 0$ with $a \neq b$, the chain of inequalities*

$$\begin{aligned} A^{2/3}G^{1/3} \quad &< \quad \sqrt{\tfrac{2}{3}AG + \tfrac{1}{3}A^2} < \sqrt{AX} < A^{1/3}\left(\tfrac{A+G}{2}\right)^{2/3} \\ &< \quad P < \frac{2\sqrt{2}\sqrt{A^2+AG}+A+G}{6} < \left(\tfrac{\sqrt{2}}{3}\sqrt{A+G}+\tfrac{1}{3}\sqrt{A}\right)^2 \\ &< \quad J < A^{-1/3}\left(\frac{\sqrt{2A}+\sqrt{A+G}}{2\sqrt{2}}\right)^{8/3} < \tfrac{2}{3}A + \tfrac{1}{3}G. \end{aligned} \tag{4.13}$$

Corollary 4 can be equivalently changed into

Proposition 5. *For $a, b > 0$ with $a \neq b$, we have*

$$\begin{aligned} \tfrac{\pi^2}{4e}P^2 \quad &< \quad A^{2-1/(\ln\pi-\ln 2)}P^{1/(\ln\pi-\ln 2)} < A^{2-1/\ln 2}\left(\tfrac{A+G}{2}\right)^{1/\ln 2} < \\ XA \quad &< \quad A^{2/3}\left(\tfrac{A+G}{2}\right)^{4/3} < P^2 < \frac{2^{10/3}}{\pi^2}A^{2/3}\left(\tfrac{A+G}{2}\right)^{4/3}, \end{aligned}$$

$$\begin{aligned} \sqrt{XA} \quad &< \quad P < \frac{2A+G}{3} < \frac{X+A}{2}, \\ X \quad &> \quad \frac{2G+A}{3} > \frac{P+G}{2}. \end{aligned}$$

The following is obtained from (3.23) and (3.25).

Proposition 6. *For $a, b > 0$ with $a \neq b$, we have*

$$\begin{aligned}\sqrt{\frac{8}{\pi e}}\frac{A+G}{2} &< \sqrt{PX} < \frac{A+G}{2}, \\ \sqrt{AG} &< \sqrt{PX} < \frac{A+G}{2}.\end{aligned}$$

Theorem 5 implies that

Proposition 7. *For $a, b > 0$ with $a \neq b$, we have*

$$\frac{\frac{e(\pi-2)}{\pi}X + P}{2} < \frac{A+G}{2} < \frac{P+X}{2} < \left(e^{-1} + \tfrac{2}{\pi}\right)\frac{A+G}{2}. \tag{4.14}$$

Corollary 6. *Let $a, b > 0$ with $a \neq b$. Then*

$$\sqrt{AG} < \sqrt{PX} < \frac{A+G}{2} < \frac{P+X}{2} < \left(e^{-1} + \frac{2}{\pi}\right)\frac{A+G}{2},$$

Theorem 6 can be restated in the following form.

Proposition 8. *Let $a, b > 0$ with $a \neq b$. Then*

(i) if $p \geq 6/5$, then the two-side inequality

$$\alpha G^p + (1-\alpha) A^p < X < \beta G^p + (1-\beta) A^p \tag{4.15}$$

holds if and only if $\alpha \geq (1 - e^{-p})$ and $\beta \leq 2/3$;

(ii) if $0 < p \leq 1$, then (4.15) holds if and only if $\alpha \geq 2/3$ and $\beta \leq (1 - e^{-p})$;

(iii) if $p < 0$, then (4.15) holds if and only if $\alpha \leq 0$ and $\beta \geq 2/3$;

(iv) the double inequality

$$M_p\left(G, A; \tfrac{2}{3}\right) < X < M_q\left(G, A; \tfrac{2}{3}\right) \tag{4.16}$$

holds if and only if $p \leq \ln 3$ and $q \geq 6/5$, where $M_p(x, y; w)$ is defined by (3.29).

References

[1] H. Alzer, Two inequalities for means, *C. R. Math. Pep. Acad. Sci. Canada* **9** (1987), 11-16.

[2] H. Alzer, Ungleichungen für Mittelwerte, *Arch. Math.* **47** (1986), 422-426.

[3] H. Alzer and S.-L. Qiu, Inequalities for means in two variables, *Arch. Math.* (Basel) **80** (2003), 201–215.

[4] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, New York 1997.

[5] M. Biernacki and J. Krzyz, On the monotonicity of certain functionals in the theory of analytic functions, *Annales Universitatis Mariae Curie-Sklodowska* **9** (1995) 135–147.

[6] C.-P. Chen and W.-S. Cheung, Sharp Cusa and Becker-Stark inequalities, *J. Inequal. Appl.* **2011** (2011): 136.

[7] Y.-M. Chu, Y.-F. Qiu, M.-K. Wang and G.-D. Wang, The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean, *J. Inequal. Appl.* **2010** (2010), Art. ID 436457, 7 pages.

[8] Y.-M. Chu, M.-K. Wang and Y.-F. Qiu, An optimal double inequality between power-type Heron and Seiffert means, *J. Inequal. Appl.* **2010** (2010), Art. ID 146945, 11 pages.

[9] Y.-M. Chu, M.-K. Wang and W.-M. Gong, Two sharp double inequalities for Seiffert mean, *J. Inequal. Appl.* **2011** (2011): 44, 7 pages.

[10] Y.-M. Chu, B.-Y. Long, W.-M. Gong and Y.-Q. Song, Sharp bounds for Seiffert and Neuman-Sándor means in terms of generalized logarithmic means, *J. Inequal. Appl.* **2013**, 2013:10.

[11] I. Costin and G. Toader, A nice separation of some Seiffert type means by power means, *Int. J. of Math. Math. Sci.* **2012** (2012), Art. ID 430692, 6 pages, doi:10.1155/2012/430692.

[12] Group of compilation, *Handbook of Mathematics*, Peoples' Education Press, Beijing, China, 1979. (Chinese)

[13] R. Klén, M. Visuri and M. Vuorinen, On Jordan type inequalities for hyperbolic functions, *J. Inequal. Appl.*, **2010** (2010), Art. ID 362548, 14 pages, doi:10.1155/2010/362548.
[14] O. Kouba, New bounds for the identric mean of two arguments, *J. Inequal. Pure Appl. Math.* **9**, 3 (2008), Art. 71, 6 pages.
[15] Y.-P. Lv, G.-D. Wang and Y.-M. Chu, A note on Jordan type inequalities for hyperbolic functions, *Appl. Math. Lett.* **25** (2012), 505-508.
[16] E. Neuman and J. Sándor, On the Schwab-Borchardt mean, *Math. Pannon.* **14** (2003), 253–266.
[17] E. Neuman and J. Sándor, On some inequalities involving trigonometric and hyperbolic functions with emphasis on the Cusa-Huygens, Wilker and Huygens inaequalities, *Math. Inequal. Appl.* **13**, 4 (2010), 715–723.
[18] E. Neuman, Inequalities for the Schwab-Borchardt mean and their applications, *J. Math. Inequal.* **5**, 4 (2011), 601–609.
[19] E. Neuman, Refinements and generalizations of certain inequalities involving trigonometric and hyperbolic functions, *Adv. Inequal. Appl.* **1**, 1 (2012), 1-11.
[20] A. O. Pittinger, Inequalities between arithmetic and logarithmic means, *Univ. Beograd Publ. Elektr. Fak. Ser. Mat. Fiz* **680** (1980), 15-18.
[21] J. Sándor, On the identric and logarithmic means, *Aequat. Math.* **40** (1990), 261–270.
[22] J. Sándor, Two sharp inequalities for trigonometric and hyperbolic functions, *Math. Inequal. Appl.* **15**, 2 (2012), 409–413.
[23] H.-J. Seiffert, Werte zwischen dem geometrischen und dem arithmetischen Mittel zweier Zahlen, *Elem. Math.* **42** (1987), 105–107.
[24] H.-J. Seiffert, Aufgabe 16, *Die Wurzel* **29** (1995) 221–222.
[25] H.-J. Seiffert, Ungleichungen für elementare Mittelwerte [Inequalities for elementary means], *Arch. Math. (Basel)* **64**, 2 (1995), 2, 129–131 (German).
[26] K. B. Stolarsky, Generalizations of the logarithmic mean, *Math. Mag.* **48**, 87–92 (1975).
[27] K. B. Stolarsky, The power and generalized logarithmic means, *Amer. Math. Monthly* **87** (1980), 545–548.
[28] M. K. Vamanamurthy and M. Vuorinen, Inequalities for means, *J. Math. Anal. Appl.* **183** (1994) 155–166.
[29] A. Witkowski, Interpolations of Schwab-Borchardt mean, *Math. Inequal. Appl.* 2012, in print.
[30] Zh.-H. Yang, ON the log-convexity of two-parameter homogeneous functions, *Math. Inequal. Appl.* **10**, 3 (2007), 499-516.
[31] Zh.-H. Yang, Refinements of Mitrinović-Cusa inequality, http://arxiv.org/pdf/1206.4911.pdf.
[32] Zh.-H. Yang, New sharp Jordan type inequalities and their applications, http://arxiv.org/pdf/1206.5502.pdf.
[33] Zh.-H. Yang, Sharp bounds for the second Seiffert mean in terms of power means, **2012**; available online at http://arxiv.org/pdf/1206.5494v1.pdf.
[34] Zh.-H. Yang, Sharp power means bounds for Neuman-Sándor mean, **2012**; available online at http://arxiv.org/abs/1208.0895.
[35] Zh.-H. Yang, The monotonicity results and sharp inequalities for some power-type means of two arguments, **2012**; available online at http://arxiv.org/pdf/1210.6478.pdf.
[36] Zh.-H. Yang, Refinements of a two-sided inequality for trigonometric functions, *J. Math. Inequal.* accepted.

POWER SUPPLY SERVICE CENTER, ZHEJIANG ELECTRIC POWER CORPORATION RESEARCH INSTITUTE, HANGZHOU CITY, ZHEJIANG PROVINCE, 310009, CHINA
E-mail address: `yzhkm@163.com`